

99-327198/27 B03 UYAB-97.11.07
UNIV ABERDEEN *WO 9924036-A1
97.11.07 97GB-023669 (99.05.20) A61K 31/445, 9/06, 31/70, 47/12,
38/13, 31/435

Topical formulations for treatment of dermatological conditions
e.g. psoriasis, or alopecia (Eng)

C99-096831 N(AL AM AT AU AZ BA BB BG BR BY CA CH CN CU
CZ DE DK EE ES FI GB GE GH GM HR HU ID IL IS JP
KE KG KP KR KZ LC LK LR LS LT LU LV MD MG MK
MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL
TJ TM TR TT UA UG US UZ VN YU ZW) R(AT BE CH
CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS
LU MC MW NL OA PT SD SE SZ UG ZW)

Addnl. Data: ORMEROD A D, WINFIELD A
98.11.05 98WO-GB03317

NOVELTY

Topical formulations for treating dermatological conditions
comprise:

- (a) macrocyclic lactone antibiotic, or immunosuppressive macrolide or
its active analog, derivative or prodrug; and
- (b) permeation modulator, in relative amounts such that when a
therapeutic amount is applied to the skin, a minimal systemic

B(12-M2F, 14-G2C, 14-N17, 14-R2) .4

effect is produced.

ACTIVITY

Dermatological; immunosuppressant; antibiotic.

USE

Used to treat dermatological conditions including psoriasis,
alopecia, eczema dermatitis, lichen planus, lupus erythematosus,
pyoderma gangrenosum, vitiligo, graft-versus-host disease, pustular
skin infections, bacterial skin infections or acne vulgaris (all claimed).
Used to treat diseases of the skin or mucosa (claimed).
Double-blind, left-right study compared the effect of topical sirolimus
in formulations comprising (A) sirolimus (8%) and vehicle (92%)
comprising capric acid (50%) ad benzyl alcohol (50%); and (B)
sirolimus (2.2%) in vehicle comprising isopropyl myristate (40%),
benzyl alcohol (10%) and capric acid (50%), in 24 patients with
chronic (over 3 month) plaque psoriasis. Patients had two, well-
matched, contralateral, comparable plaques about 50 cm² in area on
opposite sides of the body. A single target plaque was treated for the

WO 9924036-A+

last 6 weeks with lower potency formulation (B). After this, active
treatment was increased to higher potency formulation (A) for 6
weeks unless a clear improvement on one side had occurred. Hands
were washed thoroughly between the twice-daily application of test
formulation. The active formulation was applied consistently to one
plaque on one side of the body, while a control comprising vehicle
only was applied consistently to the plaque on the opposing side.
Assessments were performed at weeks 0, 2, 4 and 6 on the low
potency treatment and weeks 8, 10 and 12 on the higher dose
formulation, provided there were no laboratory signs of toxicity.
Clinical scoring was done at each attendance and areas traced at the
start and finish of treatment. Biopsies from active and control lesions
were performed at the end of treatment or at withdrawal. Lesions were
assessed fortnightly with subjective scoring on a scale of 0-8 for
erythema, thickening and scaling. Objective measures of improvement
were performed on both lesions at the end of each treatment period. In
biopsies, epidermal thickness was measured and immunoperoxidase
immunohistochemistry done using antibody Ki-78 to give a measure
of hyperproliferation in the epidermis and CD4 helper lymphocytes to
give a measure of autoimmune activity, which drives psoriasis. The
clinical response was measured on a scale of 0-24 with higher values
indicating a better result. For control treatment, clinical score was

9.1±4.8, ultrasound thickness 2.96±0.72 mm and erythema
measurement 33.1±7.7 (arbitrary units). For sirolimus-treated lesions,
clinical score was 11.2±5.8 (p=0.032), ultrasound thickness 2.99±0.6
mm (not significant) and erythema measurement 34.5±7.9 (arbitrary
units) (not significant). The results show that penetration of sirolimus
from the formulations does occur.

ADVANTAGE

Produces minimal systemic effects following topical
administration (claimed). Allows active agent to penetrate the skin,
particularly the stratum corneum, without significantly passing
through the epidermis into systemic (blood or lymph) systems. Avoids
side-effects associated with prior art systemic administration.
Formulation of sirolimus (2.2%) in vehicle comprising isopropyl
myristate (40%), benzyl alcohol (10%) and capric acid (50%) was
tested in single application experiments on three individuals with
normal skin. Venous blood samples were taken at 4, 7 and 24 hours
after application and no significant levels of sirolimus were detected
using MSGCMS. After 7 hours, biopsy samples were taken from two
of the individuals and bisected in parallel with the surface to give an
upper and lower half, roughly corresponding to the epidermis and

WO 9924036-A+1

99-327198/27

dermis. The skin was homogenized with acetonitrile and sirolimus
concentration determined by high-performance liquid
chromatography. The tissue concentrations of sirolimus in the upper
and lower skin segments 7 hours after application of capric
acid:isopropyl myristate:benzyl alcohol (50:40:10) containing 2.2%
sirolimus were (micro g/mg): 0 and 0.333, respectively, for subject A
and 1.5 and 0.5, respectively, for subject B. A graphical representation
is shown in the figure.

DESCRIPTION OF DRAWING

Graphical representation of the results of the clinical score (y)
determined after application of the sirolimus formulation (white bar)
and control (filled bar).

ADMINISTRATION


Administration is topical (claimed). For treatment of eczema
dermatitis, concentration of macrocyclic lactone antibiotic is 0.05-2 %
wt. %.

TECHNOLOGY FOCUS

Pharmaceuticals - Preferred Formulations: Formulations comprise up
to 10 weight % (a) and 1-60 weight % (b). The macrocyclic lactone
antibiotic is erythromycin, azithromycin or clarithromycin. The
immunosuppressant macrolide is sirolimus, FK506 or SDZ ASM 981.
(b) is an alkanolic acid or alkenic acid, preferably capric acid, octanoic
acid, oleic acid or acids of intermediate chain length. Permeation
modulator is used in conjunction with a solvent system, preferably an
aromatic alcohol or a biologically acceptable benzene derivative, with
or without an admixture of monoglycerides and/or fatty acid ester.
The permeation modulator comprises capric acid and the solvent
system comprises benzyl alcohol. The concentration of the solvent
system is 5-90 weight %. Formulations further comprise thickening
agent, preferably white soft paraffin, cetostearyl alcohol, yellow soft
paraffin, cetyl alcohol, stearyl alcohol, divalent carboxylic acid soaps
or carnauber (sic) wax.

WO 9924036-A+2

THIS PAGE BLANK (USPTO)



WO 9924036-A/3

THIS PAGE BLANK (USPTO)


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|--|--|--|--|
| (51) International Patent Classification ⁶ : A61K 31/445, 31/70, 38/13, 9/06, 47/12, 31/435 | | A1 | (11) International Publication Number: WO 99/24036 |
| | | | (43) International Publication Date: 20 May 1999 (20.05.99) |
| (21) International Application Number: PCT/GB98/03317 | | (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). | |
| (22) International Filing Date: 5 November 1998 (05.11.98) | | Published With international search report. | |
| (30) Priority Data: 9723669.9 7 November 1997 (07.11.97) GB | | | |
| (71) Applicant (for all designated States except US): ABERDEEN UNIVERSITY [GB/GB]; Auris Business Centre, 23 St. Machar Drive, Aberdeen AB2 1RY (GB). | | | |
| (72) Inventors; and (75) Inventors/Applicants (for US only): ORMEROD, Anthony, David [GB/GB]; 12 Kemnay Place, Aberdeen AB15 8SG (GB). WINFIELD, Arthur [GB/GB]; 42 Westholme Avenue, Aberdeen AB15 6AB (GB). | | | |
| (74) Agents: STEBBING, Peter, John, Hunter et al.; Ablett & Stebbing, Caparo House, 101-103 Baker Street, London W1M 1FD (GB). | | | |

(54) Title: SKIN PENETRATION ENHANCING COMPONENTS



(57) Abstract

The present invention relates to a topical formulation for the treatment of a dermatological condition which comprises a macrocyclic lactone antibiotic, immunosuppressive macrolide or a biologically active analogue, derivative or pro-drug thereof; characterized in that it further comprises a permeation modulator and the permeation modulator and the macrocyclic lactone or macrolide or the biologically active analogue, derivative or pro-drug thereof are present in relative amounts such that when a therapeutic amount is applied to the skin a minimal systemic effect is produced. The immunosuppressive macrolide may be sirolimus.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | | | TR | Turkey |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda |
| BY | Belarus | IS | Iceland | MW | Malawi | US | United States of America |
| CA | Canada | IT | Italy | MX | Mexico | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | Viet Nam |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NO | Norway | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | NZ | New Zealand | | |
| CM | Cameroon | | | PL | Poland | | |
| CN | China | KR | Republic of Korea | PT | Portugal | | |
| CU | Cuba | KZ | Kazakstan | RO | Romania | | |
| CZ | Czech Republic | LC | Saint Lucia | RU | Russian Federation | | |
| DE | Germany | LI | Liechtenstein | SD | Sudan | | |
| DK | Denmark | LK | Sri Lanka | SE | Sweden | | |
| EE | Estonia | LR | Liberia | SG | Singapore | | |

- 1 -

SKIN PENETRATION ENHANCING COMPONENTS

This present invention relates to an effective treatment for psoriasis and other dermatological conditions using a topically applied immunosuppressive agent. The preferred formulation does not allow the agent to appear in the blood or other circulatory system at any significant level.

Dermatological conditions can be uncomfortable and embarrassing for the patient, so an effective safe treatment is required. Some dermatological conditions are caused by an overactive immune system, examples are psoriasis, alopecia, lichen planus, lupus erythematosus, pyoderma gangrenosum, vitiligo and graft versus host disease. Others can be due to bacterial or pustular skin infections.

Dermatological conditions caused by an overactive immune system can be treated by immunosuppressive macrolides, for example sirolimus (rapamycin), FK-506 (tacrolimus) or SDZ ASM 981. Those that are caused by bacteria or are deeper skin infections, such as acne vulgaris and hidranitis suppurativa, can be treated by macrolide antibiotics, for example erythromycin, azithromycin and clarithromycin. The above agents may be applied by means of topical creams and lotions or taken orally.

Psoriasis affects 2.4% of the population and the current understanding of the pathogenesis of the disease is that it is driven initially by immunocytes. These and keratinocytes are mutually stimulated and activated through the production of cytokines, TGF α , IL-6 and IL-8 from lymphocytes. This leads to a hyperproliferative epidermis with rapid 36 hour cycling of the transient amplifying compartment of

- 2 -

keratinocytes.

FK506 is a macrolide antibiotic which shows part homology with sirolimus. Research in models has shown that it has some efficacy in the topical therapy of contact dermatitis, atopic eczema and to a lesser degree psoriasis. Cyclosporin is also known to be effective in treating a wide range of skin diseases. However the usefulness of these drugs is limited by their potential side effects resulting from systemic administration.

Other forms of treatment of dermatological conditions may include using topical steroids but these have undesirable effects such as irreversible atrophy and purpura.

15

In the treatment of the human or animal body, one of the considerations is that any medicament shall as far as possible affect only the afflicted part. It is well known that amounts of circulating drug should be kept as low as possible to avoid unwanted mutations. A problem with the topical application of medicaments to the skin for example, is that the medicament tends to penetrate the skin and establish itself in the circulating blood system. This is not what is intended in the treatment of dermatological conditions.

25

The macrocyclic lactone antibiotic rapamycin for example as disclosed in EP-A-0533433 has already been used topically to treat such skin disorders as psoriasis and dermatitis. However no attempt has been made to reduce the amount of rapamycin translocated across the skin into the systemic system. Nor is there any discussion of the reduction of the levels of circulating rapamycin or other macrolide drug at the same time as providing therapeutically effective treatment for

- 3 -

a variety of skin disorders.

We have now found that this may be achieved by the addition to such drugs of a permeation modulator. Permeation enhancers
5 are well known as a class of drug translocation facilitators, but the purpose of these is to increase the drug flux across the skin. A permeation modulator however has the facility to allow the drug to penetrate the skin, and particularly the stratum corneum, without significantly passing through the
10 epidermis into systemic systems (eg the blood or lymph systems).

It is also known that immunosuppressive agents taken orally and steroids applied topically can be used to treat
15 dermatological conditions, such as psoriasis or eczema. However, they are often non-specific in their action which leads to undesirable side effects. Thus it would be desirable to develop a topical delivery formulation for an immunosuppressive agent which preferentially treats the
20 diseased sites only and avoids significant systemic exposure; so reducing harmful side effects.

Sirolimus is a macrocyclic lactone antibiotic produced by the organism *Streptomyces hygroscopicus*; it is known to have
25 potent immunosuppressive activities. Sirolimus acts through specific binding of a family of cytosolic immunophilins called the FK binding proteins (FKBP). The sirolimus FKBP complex acts at least three sites. Firstly, by blocking the phosphorylation activation of p70 s6 kinase, an enzyme acting
30 on the 40S ribosomal subunit s6 protein, thereby reducing the efficiency of translation. Secondly by preventing activation of specific elongation factors required for protein synthesis. Thirdly, it inhibits enzyme activity of the cyclin dependent

- 4 -

kinase cdK-cyclin E complex which forms one of the tight controls of the G1/S transition in cell division by inhibiting the normal decline of the p27 cdk inhibitor which would follow IL-2 stimulation. Sirolimus has an advantage over other immunosuppressive agents in the treatment of psoriasis as it has an inhibitory effect on keratinocyte proliferation. In vitro experiments have shown that this inhibitory effect takes place at concentrations ranging from 3-10 μ g/ml. A broader range may be employed for example 1 to 20 μ g/ml, but the more efficacious range is 5-8 μ g/ml.

According to the first aspect of the invention, there is provided a topical formulation for the treatment of a dermatological condition which comprises a macrocyclic lactone antibiotic or immunosuppressive macrolide or a pharmacologically active analogue, derivative or pro-drug thereof; characterised in that it further comprises a permeation modulator and the permeation modulator and the macrocyclic lactone antibiotic, immunosuppressive macrolide or pharmacologically active analogue, derivative or pro-drug are present in relative amounts such that when a therapeutic amount is applied to the skin, a minimal systemic effect is produced.

By the term "minimal systemic effect", is meant that the amount of active principal detectable in the blood stream is preferably less than 0.3 ng/ml over 4 to 24 hours after administration, more preferably below 0.1 ng/ml over the same period.

30

Preferably the macrocyclic lactone antibiotic is selected from erythromycin, azithromycin or clarithromycin. These macrocyclic lactone antibiotics are effective for treating

- 5 -

pustular and bacterial skin infections such as acne vulgaris.

Conveniently the immunosuppressive macrolide is selected from sirolimus, FK-506 or SDZ ASM 981. Sirolimus is a favoured
5 alternative because it is also an effective antibiotic which is useful in the microbiological preservation of the formulation. The microbiological properties of sirolimus are also helpful in the treatment of scalp and flexural psoriasis, seborrhoeic dermatitis and in secondarily atopic eczema.

10

In preferred embodiments the permeation modulator may be an alkanoic or alkenic acid, preferably having 6 to 20 carbon atoms such as capric acid, octanoic acid, oleic acid or acids or such acids of intermediate chain length. The permeation
15 modulator aids the penetration of the immunosuppressive macrolide or macrocyclic antibiotic through the stratum corneum, the principle barrier to the penetration of drugs. The stratum corneum is an aggregate of the stacked, flattened skeletons of keratin filled cells interspersed with lipid
20 monolayer structures and water. The addition of the permeation modulator to the formulation results in the partial disruption of the barrier components, particularly the lipid structures. A gradient of the drug can then be produced across the stratum corneum particularly, which facilitates the
25 diffusion of the immunosuppressive macrolide or macrocyclic lactone antibiotic across the stratum corneum into the living epidermis. The relative concentrations of the macrolide or antibiotic and the permeation modulator are chosen so that only partial penetration of the skin occurs; the macrocyclic
30 lactone antibiotics or immunosuppressive macrolides reach the areas which require treatment but significant absorption of the said drugs into the systemic circulation is avoided thus reducing the likelihood of any systemic side effects.

- 6 -

Conveniently the permeation modulator is used in conjunction with a solvent system which includes an aromatic alcohol such as phenyl-alkanol or a biologically acceptable benzene derivative, with or without an admixture of monoglycerides and/or a fatty acid ester (e.g. isopropyl myristate). Other solvents used, include benzaldehyde, benzyl benzoate and acetone. The combination of solvent and permeation modulator further optimises the passage of the immunosuppressive macrolide or the macrocyclic lactone antibiotic across the stratum corneum.

Preferably, the concentration of the macrocyclic lactone antibiotic or immunosuppressive macrolide is up to 10% by weight of the formulation. More preferably the concentration of the macrocyclic lactone antibiotic or immunosuppressive macrolide is either 0.5% to 5.9% or 6% to 12% by weight. Even more preferably the concentration of the macrocyclic antibiotic or immunosuppressive macrolide is either 1 to 5% or 6 to 8% by weight. A concentration of 0.05% to 2% is most preferable in the treatment of eczema. The term "% by weight" used herein refers to the "% by weight of the final formulation".

Preferably the above ranges of macrocyclic lactone antibiotic or immunosuppressive macrolide or analogue derivative or pro-drug thereof are used in an agent comprising a permeation modulator; wherein the concentration of the permeation modulator is 0.1% to 60% by weight. More preferably the concentration of the permeation modulator is either 0.1% to 39.9% or 40% to 80% by weight. Even more preferably the concentration of the permeation modulator is either 0.1% to 19.9%, 20% to 39.9% or 40% to 60%.

- 7 -

Preferably the above ranges of macrocyclic lactone antibiotic or immunosuppressive and permeation modulator are used in a formulation in conjunction with a solvent system; wherein the concentration of the solvent system is 5% to 90% by weight.

5 More preferably the concentration of the solvent system is either 0.1% to 49.9% or 50% to 90% by weight. Even more preferably the concentration of the solvent system is either 0.1% to 19.9%, 20% to 39.9%, 40% to 69.9% or 70% to 90% by weight.

10

Preferably a thickening agent is present in the formulation. If the formulation is to be used topically, it should be of an appropriate consistency. Therefore, thickening agents such as cetostearyl alcohol or commercially available medical grade

15 white soft paraffin may be added. These can reduce the penetration of the immunosuppressive agent but they are required for effective application. The formulations of the invention are particularly suitable for treatment of conditions of the scalp.

20

In addition to the liquid and solid vehicles set forth above, the formulations of the invention may additionally include one of the following:- flavouring agents, lubricants, solubilizers, suspending agents, filler and glidants.

25

The formulation can also be dissolved or suspended in any pharmaceutically acceptable liquid carrier or vehicle such as water or a pharmaceutically acceptable oil or fat. Such a liquid carrier or vehicle can contain other pharmaceutically

30 acceptable additives such as solubilizers, emulsifier, buffers, preservatives, suspending agents, thickening agents, colouring agents, viscosity regulators, stabilizers or osmo-regulators.

- 8 -

The invention will now be described, by way of illustration only, with reference to the following examples, tables and figures accompanying the specification

5

Figure 1 is a graphical representation of the effect on the flux ($\mu\text{g/hr/cm}^2$) of sirolimus (y) through the stratum corneum by varying the capric acid and benzyl alcohol ratio, where x is the percentage of capric acid in the benzyl alcohol.

10

Figure 2 is a graphical representation of the effect on the flux ($\mu\text{g/hr/cm}^2$) of sirolimus (y) through the stratum corneum by varying the octanoic acid and benzyl alcohol ratio, where x is the percentage of octanoic acid in the benzyl alcohol.

15

Figure 3 is a graphical representation of the effect on the flux ($\mu\text{g/hr/cm}^2$) of sirolimus (y) through the stratum corneum by varying the oleic acid and benzyl alcohol ratio, where x is the percentage of oleic acid in the benzyl alcohol.

20

Figure 4 is a graphical representation of the effect on the flux ($\mu\text{g/hr/cm}^2$) of sirolimus (y) through the stratum corneum by varying the sirolimus concentration (mg/ml) (x) while keeping the capric acid to benzyl acid ratio constant.

25

Figure 5 is a graphical representation of the results of the clinical score (y) determined after application of the sirolimus formulation () and the control (::::) in Example 3.

30

Figure 6 is a graphical representation of the difference in the clinical score after application with sirolimus formulation in Example 3, where y is the number of subjects

- 9 -

in each group. A positive score (x) shows improvement with use of the active formulation.

Figures 1 to 4 were obtained by *in vitro* experimentation. The results were used to optimize the sirolimus concentration and the ratio of permeation enhancer and solvent used in *in vivo* experiments.

Example 1

10 A formulation was formed of 8% sirolimus and 92% of a vehicle of capric acid (50%) with benzyl alcohol (50%). This was tested in single application experiments on four individuals with normal skin. Venous blood samples were taken at 4, 7 and 24 hours after application and no significant levels of
15 sirolimus were detected using MSGCMS, which is able to detect sirolimus levels down to 0.1ng/ml.

In parallel, skin biopsies were taken from the individuals after 7 hours, the biopsy samples were glued to a glass slide
20 and serially sectioned horizontally into 4 layers each 0.7mm thick and extracted with acetonitrile. The results are given in Table 1.

Table 1 shows the tissue concentrations of sirolimus 7 hours
25 after application of capric acid: benzyl alcohol (50:50) containing sirolimus at 8%. The horizontal skin sections were each 0.7mm. Accordingly, for example, the section of skin designated 2 was the horizontal layer of skin 0.7-1.4mm from the surface of the skin.

- 10 -

| Section of skin 1=surface | Sirolimus concentration $\mu\text{g}/\text{mg}$ | | | |
|------------------------------|---|-------|-------|-------|
| | A | B | C | D |
| 1 | 0.059 | 0.288 | 0.301 | 0.216 |
| 2 | Not done | 0.108 | 0.144 | 0.126 |
| 3 | 0.255 | 0.173 | 0.339 | 0.256 |
| 4 | 0.239 | 0.214 | 0.370 | 0.241 |

10 Example 2

A formulation of sirolimus (2.2%) in a vehicle comprising isopropyl myristate 40%, benzyl alcohol 10% and capric acid 50% was tested in single application experiments on three individuals with normal skin. Venous blood samples were taken at 4, 7 and 24 hours after application and no significant levels of sirolimus were detected using MSGCMS.

After 7 hours biopsy samples were taken from two of the individuals. These were bisected in parallel with the surface to give an upper and lower half, roughly corresponding to the epidermis and dermis. The skin was homogenised with acetonitrile and sirolimus concentration was determined by HPLC. The results are given in Table 2

Table 2 shows the tissue concentrations of sirolimus 7 hours after application of capric acid: isopropyl myristate: benzyl alcohol (50:40:10) containing sirolimus at 2.2%.

| Level of skin segment | Sirolimus Concentration $\mu\text{g}/\text{mg}$ | |
|-----------------------|---|-----------|
| | Subject A | Subject B |
| Upper (1) | 0 | 1.5 |
| Lower (2) | 0.333 | 0.5 |

- 11 -

Example 3

A double blind, left-right comparison of the effect of applying topical sirolimus in formulations as described in Examples 1 and 2, to 24 patients with chronic (over three months) plaque psoriasis was conducted. (22 out of the 24 patients were eventually analysed.) A single target plaque was treated for the first 6 weeks with the lower potency formulation of Example 2. After this the active treatment was increased to the higher potency formulation of Example 1 for 6 weeks unless a clear improvement on one side had already occurred.

The study included adults with stable, clearly demarcated, chronic plaque psoriasis, and two, well matched, contralateral, comparable plaques about 50cm² in area on opposite sides of the body. Subjects were all aged over 18 years, were able to apply creams and had no other significant medical problems. Transaminases were not more than twice the upper limit of normal and subjects were selected to avoid those likely to have a holiday in sunlight during the 6-12 weeks of the trial.

Before the trial started, there was a two week washout period in which only bland emollients were applied to the target lesions.

Treatment was randomised and double blind. Hands were thoroughly washed between the twice daily application of the test formulations. The active formulation was applied consistently to one plaque while a control comprising only the vehicle base was applied consistently to the plaque on the opposite side. Where possible the arms or elbows were selected as target areas as cross contamination is less likely at these

- 12 -

sites.

Assessments were done at weeks 0, 2, 4 and 6 on the low potency treatment and at 8,10 and 12 on the higher dose formulation, provided there were no signs or laboratory evidence of toxicity. Clinical scoring was done at each attendance and areas traced at the start and finish of treatment. Biopsies from active and control lesions were performed at the end of treatment or at withdrawal. Biopsies were not done if an adverse event such as a reaction to the application occurred as this would influence the measures being assessed.

The lesions were also assessed at fortnightly intervals with subjective scoring on a scale of 0-8 for erythema, thickening, and scaling. Objective measures of improvement were performed on both lesions at the end of each treatment period (low and high formulations). These included pulsed A scan ultrasound measurement of lesion thickness and erythema measured with a reflectance erythema metre, both were averaged over 5 areas in each psoriatic lesion and were validated using a previous study which was performed using betamethasone as a reference.

At each visit we measured the full blood count, biochemistry, including urea, electrolytes, liver enzymes, bilirubin, calcium, magnesium, uric acid, glucose, amylase, muscle enzymes, lipids and cholesterol. Sirolimus levels were performed every 2 weeks during therapy. Samples for sirolimus levels were stored at minus 80° C and shipped to a central reference laboratory for analysis by LC/MS/MS by Wyeth Ayerst Research.

In biopsies, epidermal thickness was measured and

- 13 -

immunoperoxidase immunohistochemistry done using the following antibodies to count cells in a blinded fashion:

Thus, antibody Ki-67 was used to give a measure of hyperproliferation in the epidermis and CD4 helper lymphocytes were used to give a measure of auto-immune activity which drives psoriasis.

Cell counting in tissues was automated, using computer assisted image analysis (Seescan). Data was analysed by Student's T test for paired data and Wilcoxon's test.

Comparison of the final scores, active vs placebo achieved significance at 0.032 by T test or Wilcoxon's test 0.0457, see Table 3 and Figures 5 and 6. The erythema measurements and ultrasound recordings were not significantly different. Three of the twenty-two patients developed contact sensitivity to the topical preparations one to benzyl alcohol, one to sirolimus and one to both of these.

20

The antibody tests with Ki-67 showed a significant reduction of proliferating cells from a mean of 83/mm³ in control to 55/mm³ with Sirolimus (rapamycin) to give a significance of P=0.027 (T test). Using CD4 cells control values were 61/mm³ against 32.7/mm³ means values following rapamycin to give a significance of P=0.0026 (T-test). The T-test were unpaired due to missing samples.

30

- 14 -

Table 3 shows the clinical response to topical sirolimus. The clinical score is measured on a scale of 0-24 with higher values indicating a better result, ultrasound thickness in mm and erythema measurement in arbitrary units.

5

| | Sirolimus | | Control | | Significance |
|-------------------------|-----------|------|---------|------|--------------|
| | Mean | S.D. | Mean | S.D. | |
| Clinical Score | 11.2 | 5.8 | 9.1 | 4.8 | p=0.032 |
| 10 Ultrasound thickness | 2.99 | 0.6 | 2.96 | 0.72 | NS |
| Erythema measurement | 34.5 | 7.9 | 33.1 | 7.7 | NS |

15 These results show that penetration of sirolimus from a formulation described above does occur. It is thought that increased adsorption would occur through the scalp to effectively treat scalp psoriasis.

- 15 -

CLAIMS:

1. A topical formulation for the treatment of a dermatological condition which comprises a macrocyclic lactone antibiotic, immunosuppressive macrolide or a pharmacologically active analogue, derivative or pro-drug thereof; characterized in that it further comprises a permeation modulator and the permeation modulator and the macrocyclic lactone antibiotic or macrolide or the pharmacologically active analogue, derivative or pro-drug thereof are present in relative amounts such that when a therapeutic amount is applied to the skin a minimal systemic effect is produced.
2. A formulation according to claim 1 comprising up to 10% by weight of the macrocyclic lactone antibiotic or the immunosuppressive macrolide or analogue, derivative or pro-drug thereof; the permeation modulator being present at 1 to 60% by weight.
3. A formulation according to either claim 1 or 2 wherein the macrocyclic lactone antibiotic is selected from erythromycin, azithromycin or clarithromycin.
4. A formulation according to either claim 1 or 2 wherein the immunosuppressive macrolide is selected from sirolimus, FK506 or SDZ ASM 981.
5. A formulation according to any preceding claim wherein the permeation modulator is an alkanolic acid or alkenic acid.
6. A formulation according to claim 5 wherein the alkanolic acid or alkenic acid is selected from capric acid, octanoic acid, oleic such acid or acids of intermediate chain length.

- 16 -

7. A formulation according to any preceding claim wherein the dermatological condition is selected from psoriasis, alopecia, eczema dermatitis, lichen planus, lupus erthematosus, pyoderma gangrenosum, vitiligo, graft versus
5 host disease, pustular skin infections, bacterial skin infections or acne vulgaris.

8. A formulation according to claim 7 wherein the dermatological condition is eczema dermatitis and the
10 concentration of macrocyclic lactone antibiotic or immunosuppressive macrolide is 0.05% to 2% by weight.

9. A formulation according to any preceding claim wherein the permeation modulator is used in conjunction with a solvent
15 system.

10. A formulation according to claim 9 wherein the solvent system comprises an aromatic alcohol or a biologically acceptable benzene derivative, with or without an admixture
20 of monoglycerides and/or a fatty acid ester.

11. A formulation according to either claim 9 or 10 wherein the permeation modulator comprises capric acid and the solvent system comprises benzyl alcohol.
25

12. A formulation according to any of claims 8 to 11 wherein the concentration of the solvent system is 5% to 90% by weight.

30 13. A formulation according to any preceding claim further comprising a thickening agent.

14. A formulation according to claim 13 wherein the

- 17 -

thickening agent is selected from white soft paraffin, cetostearyl alcohol, yellow soft paraffin, cetyl alcohol, steryl alcohol, divalent carboxylic acid soaps and carnauber wax.

5

15. A topical formulation for the treatment of a dermatological condition which comprises an immunosuppressive macrolide or a pharmacologically active analogue, derivative or pro-drug thereof; characterized in that it further
10 comprises a permeation modulator; and the permeation modulator and the macrolide or the pharmacologically active analogue, derivative or pro-drug thereof are present in relative amounts such that when a therapeutic amount is applied to the skin a minimal systemic effect is produced.

15

16. A formulation according to either claim 15 wherein the immunosuppressive macrolide is selected from sirolimus, FK506 or SDZ ASM 981.

20 17. A formulation according to claim 16 wherein the immunosuppressive macrolide is sirolimus.

18. The use in the manufacture of a topical composition for the treatment of a dermatological condition of a macrocyclic
25 lactone antibiotic or an immunosuppressive macrolide or a pharmacologically acceptable analogue, derivative or pro-drug thereof characterised in that it further comprises a permeation modulator and the permeation modulator; the macrocyclic lactone antibiotic or the immunosuppressive
30 macrolide or pharmacologically acceptable analogue, derivative or pro-drug thereof being present in relative amounts such that when a therapeutic amount is applied to the skin a minimal systemic effect is produced.

- 18 -

19. The use of claim 18 wherein the macrocyclic lactone antibiotic or immunosuppressive macrolide is present at up to 10% by weight of the composition.

5 20. The use of an immunosuppressant macrolide, a macrocyclic lactone antibiotic or a pharmacologically active analogue, derivative or pro-drug thereof in the preparation of a topical formulation as claimed in any one of claims 1 to 17.

10 21. A method for the treatment of a disease of the skin or mucosa which comprises applying thereto a topical composition comprising a macrocyclic lactone antibiotic or an immunosuppressive macrolide or a pharmacologically acceptable analogue, derivative or pro-drug thereof; characterised in
15 that it further comprises a permeation modulator; and the permeation modulator, the macrocyclic lactone antibiotic or the immunosuppressive macrolide or pharmacologically acceptable analogue, derivative or pro-drug thereof is present in relative amounts such that when a therapeutic amount is
20 applied to the skin a minimal systemic effect is produced.

22. A method according to claim 21 wherein the macrocyclic lactone antibiotic or immunosuppressive macrolide is present at up to 10% by weight of the composition.

25

23. A method according to claim 21 or 22 wherein the immunosuppressive macrolide is utilized.

BEST AVAILABLE COPY